

University of Groningen

Expansion and fragment settlement of the non-native seagrass *Halophila stipulacea* in a Caribbean bay

Smulders, Fee O. H.; Vonk, J. Arie; Engel, M. Sabine; Christianen, Marjolijn J. A.

Published in:
Marine biology research

DOI:
[10.1080/17451000.2017.1333620](https://doi.org/10.1080/17451000.2017.1333620)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Smulders, F. O. H., Vonk, J. A., Engel, M. S., & Christianen, M. J. A. (2017). Expansion and fragment settlement of the non-native seagrass *Halophila stipulacea* in a Caribbean bay. *Marine biology research*, 13(9), 967-974. <https://doi.org/10.1080/17451000.2017.1333620>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Expansion and fragment settlement of the non-native seagrass *Halophila stipulacea* in a Caribbean bay

Fee O. H. Smulders, J. Arie Vonk, M. Sabine Engel & Marjolijn J. A. Christianen

To cite this article: Fee O. H. Smulders, J. Arie Vonk, M. Sabine Engel & Marjolijn J. A. Christianen (2017) Expansion and fragment settlement of the non-native seagrass *Halophila stipulacea* in a Caribbean bay, Marine Biology Research, 13:9, 967-974, DOI: 10.1080/17451000.2017.1333620

To link to this article: <https://doi.org/10.1080/17451000.2017.1333620>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 11 Aug 2017.



Submit your article to this journal [↗](#)



Article views: 1201



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)



ORIGINAL ARTICLE



Expansion and fragment settlement of the non-native seagrass *Halophila stipulacea* in a Caribbean bay

Fee O. H. Smulders^a, J. Arie Vonk^a, M. Sabine Engel^b and Marjolijn J. A. Christianen^{c,d}

^aInstitute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, the Netherlands; ^bSTINAPA Bonaire National Parks Foundation, Bonaire, Dutch Caribbean; ^cGroningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, the Netherlands; ^dInstitute of Marine Research and Ecosystem Studies, Wageningen University, Wageningen, the Netherlands

ABSTRACT

The non-native seagrass species *Halophila stipulacea* has spread throughout the Eastern Caribbean since 2002, and could potentially impact the functioning of local seagrass ecosystems. Important characteristics for invasiveness, such as dispersal, recruitment and expansion of *H. stipulacea* at a local scale, are unknown. We assessed *H. stipulacea* expansion rates within Lac Bay, Bonaire, Dutch Caribbean (7 km²), since its establishment in 2010 and tested the settlement potential of uprooted vegetative fragments of *H. stipulacea*. Using 49 fixed locations, we observed that between 2011 and 2015 the occurrence of *H. stipulacea* in the bay increased significantly from 6% to 20% while native *Thalassia testudinum* occurrence decreased significantly from 53% to 33%. Free-floating *H. stipulacea* fragments that were collected and tethered above the sediment rooted within 10 days with a settlement success rate of 100%. The growth of settled fragments was on average 0.91 shoots d⁻¹. The ongoing shift from native *T. testudinum* to introduced *H. stipulacea* dominated meadows may have important consequences for multiple Caribbean seagrass ecosystem functions. Given the large difference in size between the two seagrass species, functions such as coastal protection, habitat structure, food availability, and the stability and resilience of these systems can be altered. The next steps towards modelling future expansion of *H. stipulacea* throughout the Caribbean and beyond should include the assessment of fragment viability and dispersal distance, and the impacts of natural and anthropogenic disturbance on vegetative fragment density, dispersion and settlement by this species.

ARTICLE HISTORY

Received 1 December 2016
Accepted 18 May 2017
Published online 11 August 2017

RESPONSIBLE EDITOR

David Thielges

KEYWORDS

Dutch Caribbean; ecosystem services; exotic seagrass; *Halophila stipulacea*; range expansion; vegetative fragmentation

Introduction

The rate at which non-native species are being introduced in aquatic ecosystems is increasing worldwide (Ruiz et al. 2000; Molnar et al. 2008). Some of these introduced species can rapidly spread and often become invasive when they negatively impact the stability and biological diversity of ecosystems (Williams 2007; Willette et al. 2014). For example, in Mediterranean and Australian seagrass ecosystems, the non-native green alga *Caulerpa taxifolia* (M.Vahl) C.Agardh has rapidly invaded and locally replaced native seagrass meadows (Williams 2007). Recent trans-oceanic introductions of seagrass species are rare, with only two documented species: *Zostera japonica* Acherson & Graebner and *Halophila stipulacea* (Forsskål) Ascherson. The introduction of *Z. japonica* from Japan to the North-Eastern Pacific led to increased seagrass habitat by colonization of bare mudflats

(Posey 1988), although competition with *Z. japonica* in combination with disturbance significantly reduced native *Zostera marina* Linnaeus performance (Bando 2006). The other introduced seagrass species, *H. stipulacea*, originates from the Red Sea and the western Indian Ocean and settled in the Mediterranean Sea after the opening of the Suez Canal in 1869. From the early 2000s onwards, *H. stipulacea* spread to various islands in the Caribbean and was first reported on Bonaire in 2010 (Ruiz & Ballantine 2004; Debrot et al. 2011; Willette et al. 2014).

In the Mediterranean, the introduction of *H. stipulacea* did not negatively impact native seagrass meadows (Duarte 2002). However, this species can potentially have a large ecological impact at introduced locations due to its extensive range expansion and high tolerance to broad salinity, irradiance and temperature ranges and substrate types (Lipkin 1975;

CONTACT Fee O. H. Smulders feesmulders@gmail.com IBED-AEE, University of Amsterdam, PO Box 94248, 1090 GE, Amsterdam, the Netherlands

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Georgiou et al. 2016). These qualities can favour the species in competition with native species and enable it to spread quickly. In the Caribbean, the potential invasiveness and ecological impact are currently unknown (Willette et al. 2014). However, recent observations in Dominica showed that *H. stipulacea* can outcompete the native pioneer species *Syringodium filiforme* Kützinger, *Halodule wrightii* Ascherson and *Halophila decipiens* Ostenfeld (Willette & Ambrose 2012; Steiner & Willette 2015). Whether *H. stipulacea* can also replace *Thalassia testudinum* K.D.Koenig, the climax species that often dominates seagrass meadows in the Caribbean, remains to be studied. A potential shift from slow-growing, structurally complex *T. testudinum* meadows to fast-growing less complex meadows with *H. stipulacea* dominance could have a large impact on seagrass ecosystem services in this region and to the carrying capacity of seagrass meadows for recovering green turtle populations (*Chelonia mydas* (Linnaeus, 1758)). Besides invading existing seagrass beds, *H. stipulacea* may also locally increase the ecosystem services by colonizing areas which were previously bare. In Dominica, the total area covered by seagrass doubled from 2008 to 2013, exclusively due to *H. stipulacea* expansion (Steiner & Willette 2015).

Seagrasses can expand via sexual reproduction and vegetative growth. Long distance dispersal is facilitated by seeds and rhizomal fragments and local expansion by belowground rhizome elongation (Marbà & Duarte 1998; Kendrick et al. 2012). However, so far only male flowers and no seeds have been recorded for *H. stipulacea* in the Eastern Caribbean (Vera et al. 2014; Willette et al. 2014). Remarkable are the large amounts of fragments present in these bays, consisting of leaves, roots and rhizomes, of *H. stipulacea* (Ruiz & Ballantine 2004; Vera et al. 2014). These fragments apparently break off easily and are uprooted after disturbance such as grazing and waves (Steiner & Willette 2015). Fragments of colonizing seagrass species, i.e. *H. wrightii* and *Halophila johnsonii* N.J.Eiseman have been reported to settle and root within two weeks in mesocosms (Hall et al. 2006). So far, *in situ* studies on the potential of vegetative fragments as a dispersal mechanism for the non-native seagrass *H. stipulacea* are lacking.

In this study we aim (1) to quantify biannual changes (2011–2015) in seagrass occurrence in Lac Bay after establishment of the non-native seagrass species *H. stipulacea* and (2) to assess the colonization potential and rate of free floating *H. stipulacea* fragments *in situ*. Furthermore, we provide recommendations for future studies to model the expansion of this non-native seagrass and assess its potential invasiveness. We expect

that *H. stipulacea* increased its number of colonized locations in Lac Bay from 2011 to 2015, given the rapid expansions reported elsewhere in the Caribbean. Secondly, we expect that colonized areas will show an increase in percentage occurrence of *H. stipulacea* and potentially a decrease in occurrence of *T. testudinum*. Additionally, *H. stipulacea* fragments are expected to be highly viable, reflected in the fast settlement (rooting) and subsequent growth from established fragments.

Material and methods

Lac Bay

Colonization of *Halophila stipulacea* and fragment settlement was determined in Lac Bay (7 km², 12° 10'N, 68°15'W), a shallow tropical clear-water bay with extensive seagrass meadows situated in the south-east of Bonaire, Dutch Caribbean (Figure 1a). Seagrass meadows in the bay, which provide one of the most important foraging grounds for juvenile green turtles in the Caribbean, were dominated by *Thalassia testudinum* (Debrot et al. 2012; Stapleton et al. 2014). Mangrove vegetation surrounds the bay, with hypersaline backwaters at the north-western side (Figure 1a). Near this mangrove border, dense and ungrazed *T. testudinum* meadows occurred in water as shallow as 0.3 m. In the deeper centre of the bay, sandy patches alternated with grazed mixed-species meadows at depths up to 6 m, consisting of *T. testudinum*, *H. stipulacea*, *Syringodium filiforme* and various macroalgae (Figure 1a).

Seagrass occurrence in Lac Bay

To map the change in presence of the non-native seagrass species *H. stipulacea* and native seagrass species throughout Lac Bay, we assessed their occurrence in 2011, 2013 and 2015. Seagrass occurrence was assessed at 49 fixed sampling locations that were spaced evenly at 250 m intervals and marked with a handheld GPS (GPS 60Cx, Garmin). At each sampling location, we counted presence or absence of *T. testudinum*, *H. stipulacea* and *S. filiforme* in each of 100 equal squares within a one m² quadrant. These measurements were taken in six adjacent plots per location by two observers wearing scuba or snorkelling equipment. The six outcomes were averaged per location, resulting in an average occurrence percentage per species per location. When seagrass was present, depth was measured twice per location, using a tape measure, and averaged. Maps were made in QGIS




Figure 1. (a) Areal overview of Lac Bay, Bonaire, with an outline of the mangrove, seagrass and coral reef areas. (b) Seagrass meadow in Lac Bay dominated by *Halophila stipulacea*. (c) Set-up of the fragment settling experiment; the orange tether is indicated by the red arrow. (d) One of the experimental *H. stipulacea* fragments 6 days after settlement. The red circle indicates the original tethered fragment. The plant material outside the red circle was grown since settlement. Map (a) by Google Maps®, photos (b,c,d) by Fee Smulders.

(Quantum GIS Development Team 2016) to visualize change in species occurrence over the years. The occurrence per species per year was averaged, as well as the percentage of locations colonized by either species. Additionally, the relative increase or decrease in occurrence of *H. stipulacea* between 2011 and 2015 was calculated per location. Because the data were not normally distributed, a non-parametric Friedman test for paired samples ($P < 0.05$) was used to test for differences over the years per species with separate post-hoc Wilcoxon signed-rank tests, using a Bonferroni adjustment (IBM SPSS Statistics 22.0 for Windows).

***Halophila stipulacea* fragment settlement**

The colonization potential of detached *H. stipulacea* fragments was determined *in situ* in November 2015, in a period of intermediate wave action and wind speeds (average wind speed of $8.4 \pm 0.17 \text{ m s}^{-1}$, near the end of the hurricane season). A total of 25 rhizomal fragments was randomly collected from the water column ($< 1 \text{ m}$ depth) in an area with an established *H. stipulacea* population (Figure 1b). The

fragments were characterized by on average 8.8 shoots (range 3–15), 15.5 leaves (5–28) and 2.0 roots (0–7). The fragments were attached to a rope (W:L; 6 mm:70 cm) that served as a tether to keep the fragments in place. Five fragments per tether were attached with cable ties and spaced evenly (at least 10 cm interval) along the rope. This was repeated for a total of five tethers. We assumed that the settlement of a single fragment was not influenced by the other fragments on the tether and therefore considered each fragment to be a unique individual measurement. The tethers were placed just above ($< 2 \text{ cm}$) the sediment surface so that the fragments were still hanging in the water and could move on the currents. Each tether was attached diagonally to two iron pins in a 0.25 m^2 square patch of bare sand from which all seagrass biomass was removed (Figure 1c). This patch was located within a shallow mixed seagrass meadow consisting of *T. testudinum*, *H. stipulacea* and *S. filiforme* (Figure 1c). Every other day the fragments were checked for settlement. A small current was created by hand, and if the fragment remained in place and roots were visibly inserted in the

sediment the fragment was considered to be settled. One fragment was lost from the tether, so calculations were based on 24 fragments. After 12 days the fragments were collected and the new shoots, leaves and roots were counted and photographed (Figure 1d). The biomass produced since settlement could easily be distinguished from the material already present on the fragment at the start of the experiment. The old, large leaves lost their colouring compared with the young, green and small leaves. Old roots were lost and old rhizomes were thick and rigid, while new rhizomes and roots were flexible, thin and white. All produced biomass was subsequently dried at 70°C and weighed. The total biomass production per fragment and the production of new plant parts were calculated from the moment a fragment was settled. New shoot growth was proportional to new rhizome growth, as *H. stipulacea* is characterized by a mono-meristematic non-leaf-replacing growth form (Short & Coles 2001).

Results

Seagrass occurrence in Lac Bay

Species occurrence in Lac Bay changed significantly through time for both *Halophila stipulacea* ($\chi^2(2) = 7.475$, $P = 0.024$, Friedman test) and *Thalassia testudinum* ($\chi^2(2) = 18.721$, $P < 0.001$, Friedman test). Following Bonferroni correction ($P < 0.017$) we found that in the 49 fixed locations, occurrence per m² of the non-native seagrass *H. stipulacea* increased significantly from on average (\pm SE) 5.9% \pm 3.1 in 2011 to 20.0% \pm 5.2 in 2015 ($Z = -2.947$, $P = 0.003$, Wilcoxon signed-rank test) but not from 2011 to 2013 (12.8% \pm 4.4) nor from 2013 to 2015 ($Z = -2.080$, $P = 0.038$; $Z = 1.344$, $P = 0.179$ respectively, Wilcoxon signed-rank tests, Figure 2a,b,c). Native seagrass *T. testudinum* showed a significant decrease in occurrence per m² from 53.2% \pm 6.4 in 2011 to 33.2% \pm 6.1 in 2015 ($Z = -3.617$, $P < 0.001$, Wilcoxon signed-rank test) and from 2011 to 2013 (40.4% \pm 6.2) and 2013 to 2015 ($Z = -3.878$, $P < 0.001$; $Z = -2.416$, $P = 0.016$ respectively, Wilcoxon signed-rank tests). *Syringodium filiforme* remained at a constant low occurrence of 3.6% \pm 1.4 per m² ($\chi^2(2) = 0.429$, $P = 0.807$, Friedman test) over this period. The average increase in occurrence of *H. stipulacea* at already colonized locations was 42.0% per fixed location for 2011 to 2015 (range in changes from -15.8% to 99.0%). *Halophila stipulacea* settled first in the deeper, central area of the bay. Within four years it rapidly spread to 10 out of 49 additional locations that were situated in the shallower areas of the bay.




Figure 2. The occurrence and distribution of *Thalassia testudinum*, *Halophila stipulacea* and *Syringodium filiforme* in (a) 2011, (b) 2013 and (c) 2015 at the 49 fixed sampling locations in Lac Bay, Bonaire.

Two of these locations recently colonized by *H. stipulacea* were previously devoid of seagrass presence. The number of fixed locations at which *T. testudinum* was present remained comparable over this period, while the average occurrence of *T. testudinum* at these colonized locations decreased by 17.5% for 2011 to 2015 (range in changes from -98.7% to 100.0%) (Figure 3). The locations at which *T. testudinum* occurred had average depths of 2.1 m in 2011, 2.0 m in 2013 and 1.8 m in 2015, all ranging from 0.3 to 4.4 m. *Halophila stipulacea* was present on average at 3.4 m depth in 2011 (2.2–4.4 m), 3.1 m in 2013 (1.0–4.4 m) and 3.2 m in 2015 (1.5–4.4 m). The sampling locations directly adjacent to the mangroves




Figure 3. The average (\pm SE) percentage occurrence of each seagrass species per m^2 per year; significant differences between years are indicated with *. The table below the occurrence graph displays the percentage of fixed sampling locations ($n = 49$) that is occupied per species in each year in Lac Bay.

showed the highest percentages of *T. testudinum* occurrence, with >90% at an average depth of 1.0 m in all years.

Halophila stipulacea fragment settlement

Within a week after tethering, 50% of the *H. stipulacea* fragments were rooted. After 10 days all 24 fragments were considered rooted and settled (Figure 1d, 4a). Shoot growth, equal to rhizome growth, was on average $0.9 \text{ fragment}^{-1} \text{ d}^{-1}$. Leaf growth accounted for $1.8 \text{ fragment}^{-1} \text{ d}^{-1}$ and root growth for $0.7 \text{ fragment}^{-1} \text{ d}^{-1}$. Finally, total biomass production was $7.1 \text{ mg fragment}^{-1} \text{ d}^{-1}$. We observed a high variation in production of new shoots, leaves, roots and biomass per fragment (Figure 4b).




Figure 4. (a) The proportion of settled *Halophila stipulacea* fragments in time after tethering. (b) Production per fragment per day of the different plant parts (shoot, leaf, root) and total biomass (median and variation, $n = 24$).

Discussion

In this study we observed rapid expansion of the potentially invasive seagrass *Halophila stipulacea* between 2011 and 2015 throughout Lac Bay. *Halophila stipulacea* occurrence increased by about 350% within four years, and partially replaced or excluded native *Thalassia testudinum* within the fixed sampling locations. If these trends continue, this would result in an average occurrence of 26% for *T. testudinum* and 27% for *H. stipulacea* by 2017, hence potentially a shift in dominance of seagrass within seven years after settlement in this bay. *Halophila stipulacea* colonization and expansion was more extensive in the deeper parts of the bay. Here, *T. testudinum* and *S. filiforme* are intensively cropped by grazing green turtles, resulting in low-density *T. testudinum* meadows that may be more easily colonized by the fast-growing *H. stipulacea* than ungrazed, more structurally complex seagrass canopies. However, *H. stipulacea* expansion is not limited to these particular conditions, as more shallow sites have also recently been colonized and it has been observed to grow in between dense, ungrazed *T. testudinum* meadows at depths up to 0.2 m (Fee Smulders, personal observation in 2015).

In Lac Bay, *H. stipulacea* is increasing in abundance due to rapid clonal growth. In addition, this species is probably able to spread both inside and outside the bay, as fragments easily detach, float and are able to quickly root when close to the sediment. The *H. stipulacea* fragments were abundant, drifting both on the water surface, in the water column and near the sediment. Randomly collected fragments of different sizes and lifespans all appeared viable and after tethering the fragments started to grow new roots and leaves within 10 days. Apart from our study, clonal growth rates of *H. stipulacea* have been reported

only once from a mesocosm study in the Mediterranean (Georgiou et al. 2016). Reported maximum growth rates are one shoot every two days at the maximum studied temperature of 30°C (Georgiou et al. 2016). Given our average production of 0.9 shoots per day, Caribbean *H. stipulacea* appears to grow remarkably faster, possibly due to the high temperatures and light availability in shallow Caribbean bays (Debrot et al. 2012). This short-term rooting of vegetative *H. stipulacea* fragments from the first field experiments are in line with *ex situ* experiments where *H. wrightii* and *H. johnsonii* fragments settled on the sediment and started rooting within two weeks (Hall et al. 2006). In another experiment from the same study where settlement was prevented, individual *H. wrightii* fragments remained viable for four weeks after being fragmented, and *H. johnsonii* for eight days at most (Hall et al. 2006). During this time the fragments can travel and reach new grounds to colonize.

The dispersal method via fragments might explain the past and predicted future expansion of *H. stipulacea*. Through this efficient method, *H. stipulacea* can thrive in meadows prone to natural and human-induced disturbances such as turtle grazing, storms, damage by propellers or anchors, eutrophication, and bioturbation from shrimps or rays (Steiner & Willette 2015; van Tussenbroek et al. 2016). These factors can create open patches in native beds and stimulate the release of vegetative fragments of the non-native species with the potential to settle at another location, as observed for the invasive macroalgal species *Caulerpa taxifolia* in Mediterranean meadows (West et al. 2007, 2009). Comparably, the invasive seagrass species *Z. japonica* was also reported to positively respond to disturbance by increasing invasion rates and therefore become competitively superior over *Z. marina* in the Pacific Ocean (Bando 2006).

Expansion of *H. stipulacea* into native seagrass meadows and the potential replacement of *T. testudinum* may compromise seagrass ecosystem functioning. *Thalassia testudinum* provides a dense canopy structure to support faunal assemblages, including green turtles, and forms a thick rhizome mat to stabilize the sediment (Patriquin 1973; van Tussenbroek et al. 2006). *Halophila stipulacea* has smaller leaves than *T. testudinum* (Den Hartog 1970) and the leaf canopy of this potentially invasive species thus likely provides less shelter for fauna (Debrot et al. 2012). Even more noteworthy are differences in below-ground structure; *H. stipulacea* is not as firmly rooted in the sediment and lower belowground biomass may result in decreased coastal protection (Christianen

et al. 2013) and eventually to changed stability and resilience of the meadows (Vonk et al. 2015). Shifts in species composition as a result of seagrass invasion are therefore expected to have a major impact on productivity, habitat structure, food availability and carbon sequestration (Marbà et al. 2015; Unsworth et al. 2015). However, *H. stipulacea* has also colonized two locations that previously consisted solely of bare sediment in Lac Bay. Therefore, *H. stipulacea* can locally increase the total cover of seagrass area and enhance ecosystem functioning, especially if this species continues to colonize unvegetated areas throughout the Caribbean (Steiner & Willette 2015).

Although *H. stipulacea* has been reported to be invasive, there are no unequivocal results supporting this claim. Our first quantification of properties of introduced seagrass fragments calls for further investigation of fragment characteristics and interspecies competition in order to determine the invasiveness of *H. stipulacea* and be able to model non-native seagrass expansion. To fill in the most important gaps of knowledge around *H. stipulacea* invasiveness potential and expansion, more knowledge is needed on fragment properties (viability, floating duration, dispersal distance, uprooting factors, density, lifespan; Ceccherelli & Cinelli 1999; Smith & Walters 1999; Hall et al. 2006; Grech et al. 2016) and the impact of natural and human-induced stressors (e.g. grazing, storms, anchor or propeller damage; Bando 2006; West et al. 2007, 2009). From the parameters obtained, we can model the rate of expansion, predict changes in meadow composition, and provide management recommendations to maintain key ecosystem services of Caribbean seagrass meadows.

Acknowledgements

We would like to thank Caren Eckrich (STINAPA) for assisting in the field with the seagrass mapping, and Lisa Becking for suggestions on the research setup and logistic support.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was supported by the Volkert van der Willigen grant 2015 of the Amsterdam University Fund to Fee Smulders and NWO (Nederlandse Organisatie voor Wetenschappelijk Onderzoek) grant no. 858.14092 to Marjolijn Christianen.

ORCID

Fee O. H. Smulders  <http://orcid.org/0000-0003-4124-8355>

References

- Bando KJ. 2006. The roles of competition and disturbance in a marine invasion. *Biological Invasions* 8:755–763. doi:10.1007/s10530-005-3543-4
- Ceccherelli G, Cinelli F. 1999. The role of vegetative fragmentation in dispersal of the invasive algae *Caulerpa taxifolia* in the Mediterranean. *Marine Ecology Progress Series* 182:299–303. doi:10.3354/meps182299
- Christianen MJA, van Belzen J, Herman PMJ, van Katwijk MM, Lamers LPM, van Leent PJM, Bouma TJ. 2013. Low-canopy seagrass beds still provide important coastal protection services. *PLoS One* 8:e62413. 8 pages. doi:10.1371/journal.pone.0062413
- Debrot AO, Hylkema A, Vogelaar W, Meesters HWG, Engel MS, de León R, et al. 2012. Baseline surveys of Lac Bay benthic and fish communities, Bonaire. *Journal of Experimental Marine Biology and Ecology* 376:17–25.
- Debrot AO, van Buurt G, Vermeij MJA. 2011. Preliminary overview of the exotic and invasive marine species in the Dutch Caribbean. Report C188/11. Wageningen, the Netherlands: IMARES. 29 pages.
- Den Hartog C. 1970. *The Sea-Grasses of the World*. Amsterdam: North-Holland Publishing Company. 275 pages.
- Duarte CM. 2002. The future of seagrass meadows. *Environmental Conservation* 29:192–206. doi:10.1017/S0376892902000127
- Georgiou D, Alexandre A, Luis J, Santos R. 2016. Temperature is not a limiting factor for the expansion of *Halophila stipulacea* throughout the Mediterranean Sea. *Marine Ecology Progress Series* 544:159–167. doi:10.3354/meps11582
- Grech A, Wolter J, Coles R, McKenzie L, Rasheed M, Thomas C, et al. 2016. Spatial patterns of seagrass dispersal and settlement. *Diversity and Distributions* 22:1150–1162. doi:10.1111/ddi.12479
- Hall LM, Hanisak MD, Virnstein RW. 2006. Fragments of the seagrasses *Halodule wrightii* and *Halophila johnsonii* as potential recruits in Indian River Lagoon, Florida. *Marine Ecology Progress Series* 310:109–117. doi:10.3354/meps310109
- Kendrick GA, Waycott M, Carruthers TJB, Cambridge ML, Hovey R, Krauss SL, et al. 2012. The central role of dispersal in the maintenance and persistence of seagrass populations. *BioScience* 62:56–65. doi:10.1525/bio.2012.62.1.10
- Lipkin Y. 1975. *Halophila stipulacea*, a review of a successful immigration. *Aquatic Botany* 1:203–215. doi:10.1016/0304-3770(75)90023-6
- Marbà N, Arias-Ortiz A, Masqué P, Kendrick GA, Mazarrasa I, Bastyan GR, et al. 2015. Impact of seagrass loss and subsequent revegetation on carbon sequestration and stocks. *Journal of Ecology* 103:296–302. doi:10.1111/1365-2745.12370
- Marbà N, Duarte CM. 1998. Rhizome elongation and seagrass clonal growth. *Marine Ecology Progress Series* 174:269–280. doi:10.3354/meps174269
- Molnar JL, Gamboa RL, Revenga C, Spalding MD. 2008. Assessing the global threat of invasive species to marine biodiversity. *Frontiers in Ecology and the Environment* 6:485–492. doi:10.1890/070064
- Patriquin DG. 1973. Estimation of growth rate, production and age of the marine angiosperm *Thalassia testudinum* König. *Caribbean Journal of Science* 13:111–123.
- Posey MH. 1988. Community changes associated with the spread of an introduced seagrass, *Zostera japonica*. *Ecology* 69:974–983. doi:10.2307/1941252
- Quantum GIS Development Team. 2016. Quantum GIS Geographic information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>.
- Ruiz GM, Fofonoff PW, Carlton JT, Wonham MJ, Hines AH. 2000. Invasion of coastal marine communities in North America: apparent patterns, processes, and biases. *Annual Review of Ecology and Systematics* 31:481–531. doi:10.1146/annurev.ecolsys.31.1.481
- Ruiz H, Ballantine DL. 2004. Occurrence of the seagrass *Halophila stipulacea* in the tropical West Atlantic. *Bulletin of Marine Science* 75:131–135.
- Short FT, Coles RG. 2001. Methods for the measurement of seagrass growth and production. In: Short FT, Coles RG, editors. *Global Seagrass Research Methods*. Amsterdam: Elsevier Science, p 155–182.
- Smith CM, Walters LJ. 1999. Fragmentation as a strategy for *Caulerpa* species: fates of fragments and implications for management of an invasive weed. *Marine Ecology* 20:307–319. doi:10.1046/j.1439-0485.1999.2034079.x
- Stapleton S, Nava M, Willes S, Brabec B. 2014. Research and Monitoring of Bonaire's Sea Turtles: 2014 Technical Report. Kralendijk, Bonaire: Sea Turtle Conservation Bonaire. 22 pages.
- Steiner SCC, Willette DA. 2015. The expansion of *Halophila stipulacea* (Hydrocharitaceae, Angiospermae) is changing the seagrass landscape in the commonwealth of Dominica, Lesser Antilles. *Caribbean Naturalist* 22:1–19.
- van Tussenbroek BI, van Katwijk MM, Bouma TJ, van der Heide T, Govers LL, Leuven RSEW. 2016. Non-native seagrass *Halophila stipulacea* forms dense mats under eutrophic conditions in the Caribbean. *Journal of Sea Research* 115:1–5. doi:10.1016/j.seares.2016.05.005
- van Tussenbroek BI, Vonk JA, Stapel J, Erftemeijer PLA, Middelburg JJ, Ziemann JC. 2006. The biology of *Thalassia*: paradigms and recent advances in research. In: Larkum AWD, Orth RJ, Duarte DM, editors. *Seagrasses: Biology, Ecology and Conservation*. Dordrecht, the Netherlands: Springer, p 409–439.
- Unsworth RKF, Collier CJ, Waycott M, McKenzie LJ, Cullen-Unsworth LC. 2015. A framework for the resilience of seagrass ecosystems. *Marine Pollution Bulletin* 100:34–46. doi:10.1016/j.marpolbul.2015.08.016
- Vera B, Collado-Vides L, Moreno C, van Tussenbroek BI. 2014. *Halophila stipulacea* (Hydrocharitaceae): a recent introduction to the continental waters of Venezuela. *Caribbean Journal of Science* 48:66–70. doi:10.18475/cjos.v48i1.a11
- Vonk JA, Christianen MJA, Stapel J, O'Brien KR. 2015. What lies beneath: why knowledge of belowground biomass dynamics is crucial to effective seagrass management.

- Ecological Indicators 57:259–267. doi:10.1016/j.ecolind.2015.05.008
- West EJ, Barnes PB, Wright JT, Davis AR. 2007. Anchors aweigh: fragment generation of invasive *Caulerpa taxifolia* by boat anchors and its resistance to desiccation. Aquatic Botany 87:196–202. doi:10.1016/j.aquabot.2007.06.005
- West EJ, Davis AR, Barnes PB, Wright JT. 2009. The role of recreational activities in creating fragments of invasive *Caulerpa taxifolia*. Journal of Experimental Marine Biology and Ecology 376:17–25. doi:10.1016/j.jembe.2009.05.015
- Willette DA, Ambrose RF. 2012. Effects of the invasive seagrass *Halophila stipulacea* on the native seagrass, *Syringodium filiforme*, and associated fish and epibiota communities in the Eastern Caribbean. Aquatic Botany 103:74–82. doi:10.1016/j.aquabot.2012.06.007
- Willette DA, Chalifour J, Debrot AOD, Engel MS, Miller J, Oxenford HA, et al. 2014. Continued expansion of the trans-Atlantic invasive marine angiosperm *Halophila stipulacea* in the Eastern Caribbean. Aquatic Botany 112:98–102. doi:10.1016/j.aquabot.2013.10.001
- Williams SL. 2007. Introduced species in seagrass ecosystems: status and concerns. Journal of Experimental Marine Biology and Ecology 350:89–110. doi:10.1016/j.jembe.2007.05.032